

VERIFICATION OF TRANSLATION

I, Naoto Sato

of 6th Fl., Kantetsu Tsukuba-Science-City Bldg. 1-1-1, Oroshi-machi, Tsuchiura, Ibaraki,
JAPAN

declare as follows:

1. That I am well acquainted with both the English and Japanese languages, and
2. That the attached document is a true and correct translation of a certified copy of the following application, which was made by me to the best of my knowledge and belief of.

Japanese Patent Application No. 2002-313076

Entitled: "METHOD TO ESTIMATE THE INHERITED RESISTANCE AND
SUSCEPTIBILITY TO RNA VIRUS DISEASE IN PIGS"

Filed on October 28, 2002

September 26, 2008

Date

Naoto

Naoto Sato

【Name of Document】 APPLICATION FOR PATENT

【Identification Number】 MOA-A0214

【Filing Date】 October 28, 2002

【Addressee】 Esq. Commissioner of the Patent Office

【Inventor】

【Address】 26-405-501, Matsushiro 4chome, Tsukuba-shi,

Ibaraki

【Name】 MITSUHASHI, Tadayoshi

【Applicant】

【Identification Number】 501167644

【Name or Appellation】 National Institute of Agrobiological Sciences

【Attorney】

【Identification Number】 100102978

【Patent Attorney】

【Name or Appellation】 Hatsushi Shimizu

【Nominated Attorney】

【Identification Number】 100108774

【Patent Attorney】

【Name or Appellation】 Kazunori Hashimoto

【Payment】

【Registration Number】 041092

【Amount】 21000 yen

【List of Attached Documents】

【Name of Document】 Specification 1

【Name of Document】 Drawings 1

【Name of Document】 Abstract 1

【Proof】 Necessary

[Document Name] Specification

[Title of the Invention] Method to estimate the inherited resistance and susceptibility to RNA virus disease in pigs

[Claims]

[Claim 1] A method for determining a pig's resistance to an influenza virus, wherein the method comprises the step of detecting an 11-bp deletion in a swine Mx1 gene exon, wherein the deletion is from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1.

[Claim 2] The method according to claim 1, comprising the steps of:

- (a) preparing a DNA sample from a subject pig;
- (b) amplifying a DNA that is a swine Mx1 gene exon and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1; and
- (c) determining the nucleotide sequence of the amplified DNA.

[Claim 3] The method according to claim 1, comprising the steps of:

- (a) preparing a DNA sample from a subject pig;
- (b) digesting the prepared DNA with a restriction enzyme;
- (c) separating DNA fragments based on their size; and
- (d) comparing the sizes of detected DNA fragments with that of a control.

[Claim 4] The method according to claim 1, comprising the steps of:

- (a) preparing a DNA sample from a subject pig;
- (b) amplifying a DNA that is a swine Mx1 gene exon and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1;
- (c) digesting the amplified DNA with a restriction enzyme;
- (d) separating DNA fragments based on their size; and
- (e) comparing the sizes of detected DNA fragments with that of a control.

[Claim 5] The method according to claim 1, comprising the steps of:

- (a) preparing a DNA sample from a subject pig;
- (b) amplifying a DNA that is a swine Mx1 gene exon and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1;
- (c) dissociating the amplified DNA into single strands;
- (d) separating the dissociated single-stranded DNAs on a non-denaturing gel; and
- (e) comparing the gel mobility of the fractionated single-stranded DNAs with that of a control.

[Claim 6] The method according to claim 1, comprising the steps of:

- (a) preparing a DNA sample from a subject pig;
- (b) amplifying a DNA that is a swine Mx1 gene exon and comprises the nucleotide sequence

from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1;
(c) determining the molecular weight of the DNA amplified in step (b) by mass spectrometry;
and
(d) comparing the molecular weight determined in step (c) with that of a control.

5 [Claim 7] The method according to claim 1, comprising the steps of:

- (a) preparing a DNA sample from a subject pig;
- (b) amplifying a DNA that is a swine Mx1 gene exon and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1;
- (c) preparing a substrate with an immobilized nucleotide probe;
- 10 (d) contacting the DNA prepared in step (b) with the substrate prepared in step (c);
- (e) determining the intensity of hybridization between the DNA and the nucleotide probe immobilized on the substrate; and
- (f) comparing the intensity determined in step (e) with that of a control.

 [Claim 8] The method according to claim 1, comprising the steps of:

- 15 (a) preparing a protein sample from a subject pig; and
- (b) determining the amount of a mutant swine Mx1 protein in the protein sample, wherein said mutant swine Mx1 protein is encoded by a nucleotide sequence that is a swine Mx1 gene exon in which the 11-bp nucleotide sequence from positions 2064 to 2074 in SEQ ID NO: 1 has been deleted.

20 [Claim 9] An oligonucleotide that is a PCR primer for the method for determining of claim 1, wherein the oligonucleotide is used to amplify a DNA region that is a swine Mx1 gene exon and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1.

 [Claim 10] An oligonucleotide comprising at least 15 nucleotides, and hybridizing to a
25 DNA region that is a swine Mx1 gene exon and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1, or a DNA region that is a swine Mx1 gene exon and comprises a nucleotide sequence in which the nucleotide sequence from positions 1964 to 1974 has been deleted.

 [Claim 11] An antibody recognizing a mutant swine Mx1 protein encoded by the
30 nucleotide sequence of a swine Mx1 gene exon in which the nucleotide sequence from positions 2064 to 2074 in SEQ ID NO: 1 has been deleted.

 [Claim 12] A test reagent for determining a pig's resistance to an influenza virus, wherein the reagent comprises the oligonucleotide according to claim 9 or 10, or the antibody according to claim 11.

[Detailed Description of the Invention]

[Technical Field of Industrial Application]

[0001]

5 The present invention relates to methods for determining genetic resistance of pigs to diseases caused by RNA viruses.

[Prior Art]

[0002]

10 Mouse studies have shown that the Mx gene encodes a protein that suppresses propagation (inhibits RNA synthesis) of RNA viruses belonging to the orthomyxovirus family (single-stranded linear RNA viruses), such as influenza viruses.

[0003]

15 One of the two swine Mx genes is the Mx1 gene, consisting of 14 exons. Some pigs of the Meishan breed have a 3-bp (Ser) deletion in exon 13. On the other hand, some animals of pig breeds that are domesticated in the West and widely used around the world, such as Landrace and Duroc, have an 11-base deletion in the last exon (see Non-Patent Document 1).

[0004]

20 The present inventors have previously reported the presence of Mx1 deletion genotypes and their frequencies (see Non-Patent Document 1) through collaboration with the STAFF Institute. Among these deletion genotypes, an 11-base deletion type without virus-suppressing ability was found to exist at a high frequency in Landrace, which is the most commonly used swine breed in the world. Landrace (Fig. 1 photograph) has been widely used to raise three-way crossbreds for fattening/production.

[0005]

25 On the other hand, the 11-base deletion has not been found in wild or half-wild species such as the Japanese wild boar and Meishan pig.

[0006]

30 The homozygous C was conceivably against the survival of pigs, since, for example, the percentage of homozygous C present in the Landrace breed was lower than that of the heterozygous C, and in other breeds only heterozygous but no homozygous pigs exist. No homozygous (C/C) pigs were found in the survey of about 40 pigs of white Western breeds, including Landrace, in the old National Institute of Animal Industry in the year 2000.

[0007]

35 The largest worldwide influenza epidemic in history, called the Spanish influenza epidemic, occurred between 1917-1918 and led to 20-40 million deaths. In Japan, more than 300,000 people died. The Spanish influenza epidemic, which started in the Northwest region of

the United States, killed approximately 600,000 people in the United States alone. At that time, a cold outbreak in the pig population was also reported. There was also a report on the phylogenetic analysis of an RNA virus obtained from formalin-preserved, paraffin-embedded lung tissue samples taken from soldiers who died during the epidemic. According to the report, the influenza virus is very close to those infectious for pigs.

[0008]

Pig farming is still popular in the Northwest region of the United States including the Iowa State. There is no detailed information on pig farming at the time of the outbreak, but if the Landrace breed or its crossbreeds were being raised on a large scale, pig populations having a high percentage of pigs susceptible to influenza viruses may have been acting as a breeding ground for the new virus and resulting in the pandemic.

[0009]

Farm animals are bred at high densities within limited areas of fixed locations. High-density breeding can mean major damages to production as disease spreads easily through the herd after an initial outbreak. As is evident from examples of influenza transmission from pigs to humans, high-density livestock herds sometimes can act as a breeding ground that transmits diseases to populations across species. To prevent such risks, preventive vaccination is used nowadays, and antibiotics-containing feed is given to animals even if they are in a healthy state. In spite of such preventive measures, great losses in pig production can occur, as exemplified by the outbreak of a new type influenza virus H3N2 in the United States between 1998-2000. Routine administration of antibiotics may result in antibiotics remaining in meat products, and therefore trigger food safety concerns among consumers. However, the preventive measures are necessary for reducing livestock losses from diseases. Genetic information governing potential disease resistance in livestock is required to maintain food safety while retaining or improving the current rate of disease suppression. Nevertheless, such information is still yet to be found.

[0010]

In addition, so far there is no effective method available for determining genetic resistance of pigs to diseases caused by RNA viruses.

[0011]

[Non-Patent Document 1]

Morozumi T. *et al.*, "Three types of polymorphisms in exon 14 in porcine Mx1 gene", *Biochemical Genetics.*, (2001) Vol. 39, p. 251-260

[Problems to be Solved by the Invention]

[0012]

The present invention was achieved under the above circumstances. An objective of the invention is to provide methods for determining genetic resistance of pigs to diseases caused by RNA viruses.

5 [Means for Solving the Problems]

[0013]

Some particular livestock and livestock breeds may have the genetically-inherited disease resistance characteristic of "low productivity, but high resistance to certain diseases". If such a hereditary characteristic of disease resistance can be introduced into breeds with high productivity, breeds of high productivity and high disease resistance can be created. This would enable the reduction of the amounts of antibiotics required. It has been difficult to identify the genetic basis governing such hereditary characteristics at the molecular level. However, the advancement of molecular genetics has accelerated the elucidation of genetic bases of various hereditary characteristics in human and mouse. If the resistance to a particular disease can be related to some of the genetic information in farm animals, disease resistant livestock can be selected and bred based on such genetic information.

[0014]

In a previous study, the inventors revealed that among domesticated pigs, some have an 11-base deletion in the gene encoding the Mx1 protein, which is responsible for the suppression of myxovirus propagation (already disclosed in a published document). Then, the inventors noticed a lower ratio of animals with homozygous deletion genotype compared to those with heterozygous deletion genotype. The inventors understood from pig farming experiences that pigs have poor resistance to respiratory diseases and that this has a huge impact on piglet production. The present inventors then investigated the effects of the 11-bp deletion in the Mx1 gene on the ability to suppress propagation of influenza viruses (members of the myxovirus family), and revealed that the 11-bp deletants completely lost the ability to suppress virus propagation, and thereby completed the present invention.

[0015]

More specifically, in Mx1 genes carrying the 11-base deletion in the last exon, the 11-base deletion causes frame shift of codons (single units of triplicate nucleotides), which relocates the stop codon to a much further position downstream and thus dramatically alters the downstream amino acid sequence. The resultant mutant Mx1 protein has a different molecular weight and structure compared to the normal (wild-type) protein, and has probably lost its ability to suppress virus propagation because of that.

[0016]

Next, the present inventors introduced a normal (wild-type) Mx1 gene, a mutant Mx1

gene containing a 3-base (3-bp) deletion in exon 13, a mutant Mx1 gene containing the 11-base (11-bp) deletion in the last exon, or an empty vector into murine 3T3 cells which have no Mx1 activity, and performed influenza A virus infection experiments.

[0017]

5 The results showed that the 11-bp deletant completely lost its ability to suppress virus propagation (arrow on the far left of Fig. 2), and had a virus propagation comparable to that of the empty vector, which was 10-100 times higher than the wild type. In addition, the 3-bp deletant had a virus propagation curve comparable to that of the wild type and the deletion did not affect the virus suppression ability. The deletant thus maintained the ability to suppress
10 virus propagation (arrow on the far right of Fig. 2). Similar results were obtained in a separate experiment of virus infection at a multiplicity of infection that is one order of magnitude smaller (MOI = 1).

[0018]

15 In particular, among the domestic pigs used around the world, there are some that carry an Mx1 gene that lacks the ability to suppress viral propagation. According to the experimental results of the present invention, such pigs are expected to have a serious defect in their defense ability during the early phase defense against an RNA virus invasion.

[0019]

20 Specifically, the above-described genotype of the swine Mx1 gene can be used to determine the genetic resistance of pigs to diseases caused by RNA viruses. In addition, it is important to eliminate the 11-base Mx1 deletants not only for the health of livestock, but also for eliminating threats that arise from emergence of new influenza viruses that are infectious to humans.

[0020]

25 The present invention relates to methods for determining genetic resistance of pigs to diseases caused by RNA viruses. More specifically, the present invention provides:

[1] a method for determining a pig's resistance to an influenza virus, wherein the method comprises the step of detecting an 11-bp deletion in a swine Mx1 gene exon, wherein the deletion is from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1;

30 [2] the method according to [1], comprising the steps of:

(a) preparing a DNA sample from a subject pig;

(b) amplifying a DNA that is a swine Mx1 gene exon and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1; and

(c) determining the nucleotide sequence of the amplified DNA;

35 [3] the method according to [1], comprising the steps of:

(a) preparing a DNA sample from a subject pig;

- (b) digesting the prepared DNA with a restriction enzyme;
- (c) separating DNA fragments based on their size; and
- (d) comparing the sizes of detected DNA fragments with that of a control;

[4] the method according to [1], comprising the steps of:

- (a) preparing a DNA sample from a subject pig;
- (b) amplifying a DNA that is a swine Mx1 gene exon and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1;
- (c) digesting the amplified DNA with a restriction enzyme;
- (d) separating DNA fragments based on their size; and
- (e) comparing the sizes of detected DNA fragments with that of a control;

[5] the method according to [1], comprising the steps of:

- (a) preparing a DNA sample from a subject pig;
- (b) amplifying a DNA that is a swine Mx1 gene exon and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1;
- (c) dissociating the amplified DNA into single strands;
- (d) separating the dissociated single-stranded DNAs on a non-denaturing gel; and
- (e) comparing the gel mobility of the fractionated single-stranded DNAs with that of a control;

[6] the method according to [1], comprising the steps of:

- (a) preparing a DNA sample from a subject pig;
- (b) amplifying a DNA that is a swine Mx1 gene exon and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1;
- (c) determining the molecular weight of the DNA amplified in step (b) by mass spectrometry; and

- (d) comparing the molecular weight determined in step (c) with that of a control;

[7] the method according to [1], comprising the steps of:

- (a) preparing a DNA sample from a subject pig;
- (b) amplifying a DNA that is a swine Mx1 gene exon and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1;
- (c) preparing a substrate with an immobilized nucleotide probe;
- (d) contacting the DNA prepared in step (b) with the substrate prepared in step (c);
- (e) determining the intensity of hybridization between the DNA and the nucleotide probe immobilized on the substrate; and

- (f) comparing the intensity determined in step (e) with that of a control;

[8] the method according to [1], comprising the steps of:

- (a) preparing a protein sample from a subject pig; and

(b) determining the amount of a mutant swine Mx1 protein in the protein sample, wherein said mutant swine Mx1 protein is encoded by a nucleotide sequence that is a swine Mx1 gene exon in which the 11-bp nucleotide sequence from positions 2064 to 2074 in SEQ ID NO: 1 has been deleted;

[9] an oligonucleotide that is a PCR primer for the method for determining of [1], wherein the oligonucleotide is used to amplify a DNA region that is a swine Mx1 gene exon and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1;

[10] an oligonucleotide comprising at least 15 nucleotides, and hybridizing to a DNA region that is a swine Mx1 gene exon and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1, or a DNA region that is a swine Mx1 gene exon and comprises a nucleotide sequence in which the nucleotide sequence from positions 1964 to 1974 has been deleted;

[11] an antibody recognizing a mutant swine Mx1 protein encoded by the nucleotide sequence of a swine Mx1 gene exon in which the nucleotide sequence from positions 2064 to 2074 in SEQ ID NO: 1 has been deleted;

[12] a test reagent for determining a pig's resistance to an influenza virus, wherein the reagent comprises the oligonucleotide according to [9] or [10], or the antibody according to [11].

[Mode for Carrying Out the Invention]

[0021]

The present invention provides methods for determining pig resistance to an influenza virus, comprising the step of detecting an 11-bp deletion in an exon of the swine Mx1 gene, wherein the deletion is from position 2064 to position 2074 in the nucleotide sequence of SEQ ID NO: 1.

[0022]

The nucleotide sequence of the last exon of the Mx1 gene described above is shown in SEQ ID NO: 1, and the amino acid sequence of a protein encoded by the nucleotide sequence is shown in SEQ ID NO: 2. Those skilled in the art can readily obtain this sequence information using the PubMed Accession No. M65087.

[0023]

The "deleted" 11-bp of the present invention has the sequence of 5'-gg cgc cgg ctc-3' in the last exon of the Mx1 gene, and corresponds to the nucleotide region from positions 2064 to 2074 in SEQ ID NO: 1.

[0024]

In the present invention, a subject pig in which a heterozygous "11-bp deletion" as

described above has been detected in the last exon of its Mx1 gene, is judged to have an Mx1 protein with no ability to suppress influenza viruses. Alternatively, the cells of a subject pig carrying a homozygous "11-bp deletion" as described above are judged to have only an Mx1 protein with no ability to suppress influenza viruses, and thus the subject pig is judged to exhibit high susceptibility to influenza viruses. A subject pig is judged to be resistant to influenza viruses if the "11-bp deletion" described above is not detected.

[0025]

In the present invention, there are no limitations on the methods (means) for detecting the above-described "11-bp deletion" in the last exon of the swine Mx1 gene, as long as the methods (means) can detect a "deletion" as defined above. For example, such a method may comprise the step of directly determining the nucleotide sequence of the last exon containing the above-defined deletion portion in the Mx1 gene of a subject pig.

[0026]

In this method, first, a DNA sample is prepared from a subject pig. The DNA sample can be prepared from, for example, chromosomal DNA extracted from organs, tissues, cells, blood, oral mucosal membranes, skin, or hair of the subject pig, and alternatively cDNA or mRNA that contains no introns.

[0027]

Next step in the method involves isolating a DNA that is an exon of the swine Mx1 gene and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1. The DNA can be isolated by PCR or such using chromosomal DNA or RNA as a template and using primers that hybridize to the last exon of the swine Mx1 gene. Then, the nucleotide sequence of the isolated DNA is determined. The nucleotide sequence of the isolated DNA can be determined by using methods known to those skilled in the art.

[0028]

In the next step of the method, the determined nucleotide sequence of the DNA is compared to that of a control. In this method, the control refers to the sequence of the normal (wild type) swine Mx1 gene (for example, SEQ ID NO: 1).

[0029]

Besides the methods which directly determine the nucleotide sequence of DNA derived from a subject pig as described above, different possible methods to determine the above-described "deletion" can also be used for the determination method of the present invention.

[0030]

For example, detection of the above-described "deletion" of the present invention can also be achieved using the method described below.

[0031]

First, a DNA sample is prepared from a subject pig. Then, the prepared DNA sample is digested with restriction enzymes. In the next step, DNA fragments are fractionated based on their size. The sizes of the detected DNA fragments are then compared with that of the control.

5 In another embodiment, first, a DNA sample is prepared from a subject pig, followed by the amplification of a DNA that is an exon of the swine Mx1 gene and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1. The amplified DNA is then digested with restriction enzymes. In the next step, DNA fragments are fractionated based on their size. The sizes of the detected DNA fragments are then compared
10 with that of the control.

[0032]

Such methods include, for example, methods using Restriction Fragment Length Polymorphism (RFLP) and PCR-RFLP. Specifically, when a restriction enzyme recognition site is mutated (deleted), the fragment sizes yielded by restriction enzyme treatments are

15 different from the control. The region containing the mutation is amplified by PCR, and digested with different restriction enzymes. The mutation can be detected based on the difference of band mobility in electrophoresis. Alternatively, chromosomal DNA is digested with these restriction enzymes and electrophoresed. The mutation can be detected by Southern blotting using a probe DNA of the present invention. Such restriction enzymes can be
20 appropriately selected depending on the mutation. In this method, instead of genomic DNA, cDNA prepared from the RNA of a subject pig using reverse transcriptase can also be used. The cDNA can be directly digested with restriction enzymes and then analyzed by Southern blotting. Alternatively, the DNA which is an exon of the swine Mx1 gene and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1, is
25 amplified by PCR using the cDNA as a template, subjected to restriction enzyme treatments, and examined for differences in band mobility.

[0033]

In an alternative method, first, a DNA sample is prepared from a subject pig. Then, the DNA which is an exon of the swine Mx1 gene and comprises the nucleotide sequence from
30 positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1, is amplified. Then, the amplified DNA is dissociated into single-stranded DNAs. In the next step, the dissociated single-strand DNAs are fractionated on a non-denaturing gel. The electrophoretic mobility of the fractionated single-stranded DNAs is compared to that of the control.

[0034]

35 This method is exemplified by PCR-SSCP (single-strand conformation polymorphism; Cloning and polymerase chain reaction-single-strand conformation polymorphism analysis of

anonymous Alu repeats on chromosome 11. *Genomics*. 1992 Jan 1, 12(1), 139-146; Detection of p53 gene mutations in human brain tumors by single-strand conformation polymorphism analysis of polymerase chain reaction products. *Oncogene*. 1991 Aug 1, 6(8), 1313-1318; Multiple fluorescence-based PCR-SSCP analysis with postlabeling. *PCR Methods Appl.* 1995 Apr 1, 4(5), 275-282). This method has advantages such as relatively simple operation and small amounts of sample required, and is thus particularly suitable for screening large quantities of DNA samples. The principle of the method is as follows. When a double-stranded DNA fragment is dissociated into single strands, each strand forms its own conformation depending on the nucleotide sequence. The dissociated DNA stands are electrophoresed on a polyacrylamide gel containing no denaturant. The complementary single-stranded DNAs which are identical in length move to different locations because of their conformational differences. Even with a single nucleotide substitution, the single-stranded DNA conformation is altered, resulting in different electrophoretic mobility in polyacrylamide gel electrophoresis. Thus, mutations such as deletions in a DNA fragment can be detected by determining the electrophoretic mobility.

[0035]

Specifically, first, the DNA which is an exon of the swine Mx1 gene and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1, is amplified by PCR or the like. Generally, a preferred size of the PCR-amplified region ranges from about 200 to 400 bp. Those skilled in the art can appropriately select the PCR conditions and such. The amplified DNA products can be labeled by using PCR primers labeled with an isotope such as ^{32}P , fluorescent dye, biotin, or such. Alternatively, the amplified DNA products can be labeled using a PCR solution containing substrate nucleotides labeled with an isotope such as ^{32}P , fluorescent dye, biotin, or such. Alternatively, the amplified DNA fragments can be labeled by using the Klenow enzyme or such to add substrate nucleotides labeled with an isotope such as ^{32}P , fluorescent dye, biotin, or such after PCR is completed. The labeled DNA fragments are denatured by heating, and then electrophoresed on a polyacrylamide gel containing no denaturant such as urea. The conditions for separating the DNA fragments can be improved by adding an adequate amount of glycerol (about 5-10%) to the polyacrylamide gel. The electrophoresis conditions are appropriately selected depending on the properties of the respective DNA fragments. Typically, electrophoresis is carried out at room temperature (20-25°C). When the separation is insufficient, a temperature that gives optimal mobility is selected from 4 to 30°C. Following electrophoresis, the DNA fragments are analyzed for their mobility by autoradiography using X-ray film, fluorescence scanner, or the like. Bands that have differential mobility are identified, excised directly from the gel, amplified again by PCR and then sequenced directly. The presence of a mutation can be confirmed based on the sequence. Unlabeled DNA can also be used and detected as bands by staining the gel using

ethidium bromide, silver staining, or the like, after electrophoresis is completed.

[0036]

In an alternative method, first, a DNA that is an exon of the swine Mx1 gene and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1 is prepared from a subject pig, and a substrate is prepared to which nucleotide probes that hybridize to the DNA have been immobilized. Then, the DNA is contacted with the substrate. The "deletion" described above is detected by detecting the DNAs hybridized to the nucleotide probes immobilized on the substrate.

[0037]

Such a method is exemplified by DNA array. Methods known to those skilled in the art can be used to prepare from a subject pig, a DNA sample that is an exon of the swine Mx1 gene and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1. In a preferred embodiment, the DNA sample may be prepared from, for example, chromosomal DNA extracted from tissues or cells of skin, oral mucosal membranes, blood, or the like of a subject pig. The DNA sample to be used in the method described above can be prepared from chromosomal DNA as follows. A DNA that is a swine Mx1 gene exon and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1, is prepared by, for example, PCR using chromosomal DNA as a template, and primers that hybridize to the DNA that is a swine Mx1 gene exon and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1. The prepared DNA sample can be labeled for the convenience of detection using methods known to those skilled in the art, as necessary.

[0038]

Herein, the term, "substrate" refers to a plate-shaped material onto which nucleotides can be immobilized. Herein, the term "nucleotide" refers to both oligonucleotides and polynucleotides. There are no limitations on the substrate of the present invention, as long as nucleotides can be immobilized on it. Substrates routinely used in the DNA array technology can be used suitably in the present invention.

[0039]

Typical DNA array is composed of thousands of nucleotides densely printed on a substrate. Typically, these DNAs are printed on the surface of a non-porous substrate. The surface of a typical substrate is made of glass. However, it may be made of a porous membrane, for example, nitrocellulose membrane.

[0040]

Methods of immobilizing nucleotides (array) used in the present invention include the oligonucleotide-based array method developed by Affymetrix. Oligonucleotides used in the

oligonucleotide-based array are typically synthesized *in situ*. Any of the known methods of synthesizing oligonucleotides *in situ* based on, for example, the photolithography technology (Affymetrix) or ink-jet printing technology for immobilization of substances (Rosetta Inpharmatics), can be used to prepare the substrates of the present invention.

5 [0041]

There are no limitations on the nucleotide probe to be immobilized onto a substrate, as long as it can detect the above-described "deletion". Specifically, the probe includes, for example, probes hybridizing specifically to a DNA that is a wild-type swine Mx1 gene exon and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of
10 SEQ ID NO: 1, or a DNA that is a swine Mx1 gene exon in which the nucleotide sequence from positions 2064 to 2074 of the nucleotide sequence in SEQ ID NO: 1 have been deleted. As long as specific hybridization is possible, the nucleotide probe needs not be completely complementary to the DNA to be detected which is a wild-type swine Mx1 gene exon and comprises the nucleotide sequence from positions 2064-2074 in the nucleotide sequence of SEQ
15 ID NO: 1, or the DNA which is a swine Mx1 gene exon in which the nucleotide sequence from positions 2064 to 2074 of the nucleotide sequence in SEQ ID NO: 1 have been deleted.

[0042]

In the present invention, when oligonucleotides are immobilized, the length of a nucleotide probe to be bound to a substrate is typically 10-100 bases, preferably 10-50 bases,
20 more preferably 15-25 bases.

[0043]

Next step in the present invention involves contacting a cDNA sample with the substrate. In this step, a DNA sample is hybridized to the above-described nucleotide probes. Hybridization solution and conditions vary depending on factors such as the length of the
25 nucleotide probes to be immobilized onto the substrate; however, the hybridization can generally be carried out using methods known to those skilled in the art.

[0044]

Next step in the present invention involves determining whether the DNA sample is hybridized to the nucleotide probes immobilized on the substrate, and the intensity of the
30 hybridization. This detection can be achieved by, for example, reading fluorescence signals using a scanner or such. In DNA array technology, DNA immobilized on a glass slide is generally referred to as the "probe", while labeled DNA in the solution is referred to as the "target". Therefore, the above-described nucleotides immobilized on the substrate are referred to as "nucleotide probes" herein.

35 [0045]

In addition to the methods described above, the allele specific oligonucleotide (ASO)

hybridization method can be used to detect deletions at a particular position. An oligonucleotide comprising a nucleotide sequence with potential mutations (deletions) is prepared, and hybridized with a sample DNA. Mutations in the DNA would result in a decrease in the efficiency of hybrid formation. The degree of hybridization can be determined by, for example, Southern blotting or methods using quenching resulting from the intercalation of a specific fluorescent reagent into the gaps of the hybrid. The ribonuclease A mismatch cleavage method can also be used. Specifically, a DNA that is a swine Mx1 gene exon and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1, is amplified by PCR or such. The amplified DNA is hybridized with a labeled RNA prepared from, for example, a cDNA of the last exon of the Mx1 gene incorporated into a plasmid vector. The hybrid has a partially single-stranded structure where the mutation is present. The single-stranded portion is cleaved by ribonuclease A, and detected using autoradiography or such. The mutation can be detected using the method described above.

[0046]

In an alternative method, first, DNA is prepared from a subject pig, and then a DNA which is a swine Mx1 gene exon and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1, is amplified. The amplified DNA is then analyzed by mass spectrometry to determine its molecular weight. The determined molecular weight is compared to that of a control. The method includes, for example, MALDI-TOF MS (Trends Biotechnol, 18, 77-84 (2000)).

[0047]

In another embodiment of the present invention, the above-described detection method is a detection method using the expression products of the last exon of the swine Mx1 gene as an indicator. Herein, the term "expression" refers to both transcription and translation.

Accordingly, the "expression products" include both mRNA and proteins.

[0048]

The present invention provides a method of assessing pigs that have resistance against influenza viruses, wherein the method comprises detecting expression products of the last exon of the swine Mx1 gene. In a preferred embodiment of the method, first, a protein sample is prepared from a subject pig, and then the amount of a mutant swine Mx1 protein contained in the sample is determined, wherein the mutant protein is encoded by the nucleotide sequence that is a swine Mx1 gene exon in which the 11-bp segment from positions 2064 to 2074 in SEQ ID NO: 1 has been deleted.

[0049]

Such a method includes SDS-polyacrylamide gel electrophoresis; and Western blotting, dot blotting, immunoprecipitation, enzyme-linked immunosorbent assay (ELISA), and

immunofluorescence method which use an antibody against a mutant swine Mx1 protein encoded by a nucleotide sequence that is a swine Mx1 gene exon in which the 11-bp segment from positions 2064 to 2074 in SEQ ID NO: 1 has been deleted.

[0050]

A subject pig is determined to be susceptible to influenza viruses, when a mutant swine Mx1 protein encoded by the nucleotide sequence that is a swine Mx1 gene exon in which the 11-bp segment from positions 2064 to 2074 in SEQ ID NO: 1 has been deleted, is detected using the method described above. In contrast, when the above-described mutant protein is not detected, a subject pig is determined to be resistant to influenza viruses.

[0051]

The detection method is not restricted to the various detection methods described above.

[0052]

The present invention also provides test reagents to be used in the methods of the present invention for assessing pigs that have resistance to influenza viruses.

[0053]

In an embodiment, the reagent is a test reagent comprising an oligonucleotide of at least 15 nucleotides that hybridizes to a DNA region, which is a swine Mx1 gene exon and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1, or a DNA region which is a swine Mx1 gene exon and comprises a nucleotide sequence containing a deletion of positions 1964 to 1974.

[0054]

The oligonucleotide hybridizes specifically to a DNA region that is a swine Mx1 gene exon and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1, or a DNA region that is a swine Mx1 gene exon and comprises a nucleotide sequence containing a deletion at positions 1964 to 1974. Herein, the phrase "hybridizes specifically" means that there is no significant cross-hybridization between the oligonucleotide and DNAs encoding other proteins under standard hybridization conditions or preferably under stringent hybridization conditions (for example, the conditions described in Sambrook *et al.*, "Molecular Cloning", 2nd Ed., Cold Spring Harbor Laboratory Press, New York, USA, 1989). It is not necessary for an oligonucleotide to be completely complementary to the above-described nucleotide sequence to be detected, as long as it can hybridize specifically.

[0055]

The oligonucleotide can be used as a probe or primer in the detection methods of the present invention. When the oligonucleotide is used as a primer, its length is typically 15-100 bp, and preferably 17-30 bp. There are no limitations on the primer, as long as it allows the

amplification of a DNA region that is a swine Mx1 gene exon and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1. Such primers include, though not limited to, for example, oligonucleotides comprising the sequence of SEQ ID NO: 9 or 10 described below in the Example.

5 [0056]

Alternatively, when the oligonucleotide of the present invention is used as a probe, there are no limitations on the probe, as long as it hybridizes specifically to at least a portion of a DNA region that is a swine Mx1 gene exon and comprises the nucleotide sequence from positions 2064 to 2074 in the nucleotide sequence of SEQ ID NO: 1, or a portion of a DNA region that is a swine Mx1 gene exon and comprises a nucleotide sequence containing a deletion of positions 1964 to 1974. The probe can be a synthetic oligonucleotide, and is typically at least 15 bp long.

10 [0057]

The oligonucleotides of the present invention can be prepared by, for example, using a commercially available oligonucleotide synthesizer. The probes can also be prepared as double-stranded DNA fragments from restriction enzyme treatments or the like.

15 [0058]

The oligonucleotides of the present invention can be suitably used as probes after proper labeling. The labeling method includes, for example, labeling methods that phosphorylate the 5' end of an oligonucleotide with ³²P using the T4 polynucleotide kinase; methods such as the random primer method, in which substrate nucleotides labeled with an isotope such as ³²P, fluorescent dye, biotin, or such are incorporated into DNA, by using a primer such as random hexamer oligonucleotide or the like, and a DNA polymerase such as the Klenow enzyme.

20 [0059]

In another embodiment, the test reagent of the present invention is a reagent comprising an antibody against a mutant swine Mx1 protein encoded by the nucleotide sequence that is a swine Mx1 gene exon in which the 11-bp segment from positions 2064 to 2074 in SEQ ID NO: 1 has been deleted. There are no limitations on the antibodies of the present invention, as long as they can be applied to the methods of this invention. Such an antibody includes, for example, polyclonal and monoclonal antibodies. The antibody may be labeled as necessary.

25 [0060]

The antibodies against a mutant swine Mx1 protein encoded by the nucleotide sequence that is a swine Mx1 gene exon in which the 11-bp segment from positions 2064 to 2074 in SEQ ID NO: 1 has been deleted, can be prepared by methods known to those skilled in the art. Polyclonal antibodies can be prepared by the procedure described below. Small animals such as rabbits are immunized with a mutant swine Mx1 protein, or a recombinant mutant swine Mx1 protein or a partial peptide thereof, obtained by expressing its GST fusion protein in

microorganisms such as *E. coli*. Sera collected from the animals are purified, for example, by using ammonium sulfate precipitation, protein A column, protein G column, DEAE ion exchange chromatography, or affinity column coupled with the mutant swine Mx1 protein or synthetic peptide. Monoclonal antibodies can be prepared by, for example, immunizing a small animal
5 such as mouse with a mutant swine Mx1 protein or a partial peptide thereof, extracting and gently crushing the mouse spleen to isolate cells. The cells obtained are fused with mouse myeloma cells using reagents such as polyethylene glycol. From these fused cells (hybridoma), clones producing antibodies that bind to the mutant swine Mx1 protein are selected. Next, the resulting hybridomas are transplanted into the peritoneal cavity of a mouse, from which ascites is
10 collected. The monoclonal antibody prepared is purified by, for example, using ammonium sulfate precipitation, protein A column, protein G column, DEAE ion exchange chromatography, affinity column coupled with the mutant swine Mx1 protein or synthetic peptide.

[0061]

The test reagent described above may contain, in addition to an oligonucleotide or
15 antibody as the active ingredient, sterilized water, physiological saline, a vegetable oil, a detergent, a lipid, a solubilizing agent, a buffer, a protein stabilizer (BSA, gelatin, etc.), a preservative, or such, as necessary.

[Examples]

[0062]

Herein below, the present invention will be specifically described using Examples,
however, it is not to be construed as being limited thereto.

[0063]

It was examined whether or not the Mx1 gene mutations identified in the swine breeds
25 Meishan and Landrace had an impact on the function of the expressed protein.

[0064]

Specifically, whether or not the Mx1 gene mutations affect the defense function against
infection was examined by carrying out experiments to infect cells harboring a forced-expressed
normal or mutant Mx1 gene with an influenza virus.

[0065]

[Example 1] cDNA cloning of normal and mutant Mx1 genes from Meishan and Landrace pigs

5 ml of EDTA blood was collected from each of the 39 Meishan pigs and 36 Landrace
pigs. The blood samples were diluted twice with RPMI 1640 (GIBCO), overlaid on
Ficoll-Paque (Pharmacia Biotech), and then centrifuged at 400x g for 40 minutes. The layer
35 containing lymphocytes was collected, and the cells were washed three times with phosphate
buffered saline. The prepared lymphocytes were treated with DNazol Reagent (GIBCO) for

DNA purification. PCR amplification was performed using the purified DNAs as a templates at conditions of 5 minutes at 94°C; 40 cycles of 30 seconds at 94°C, 30 seconds at 55°C, and 1 minute at 72°C; and 5 minutes at 72°C. Exon 13 was amplified from the DNA from Meishan pigs using a primer pair: 5'-CTGAAAGATCTCGGCTATGGAGG-3'/SEQ ID NO: 3; and

5 5'-AAGAAGCTGAGACGTCGATCCGGCT-3'/SEQ ID NO: 4. The last exon was amplified from the DNA from Landrace pigs using the primer pair 5'-AAGCGCATCTCCAGCCACATC-3'/SEQ ID NO: 5; and 5'-AAGACATTGGGCGTGAAAGG-3'/SEQ ID NO: 6.

The presence of the mutation in exon 13 was determined by direct sequencing. The presence of the mutation in the last exon was determined by examining the RFLP after Nal I treatment at 37°C for 2 hours.

[0066]

20 ml of EDTA blood was collected from animals that had been identified to carry a normal or mutant Mx1 gene. Lymphocytes were prepared from the blood, and suspended in RPMI 1640 containing 10% fetal bovine serum (FBS). The cell suspensions were incubated in a CO₂ incubator at 37°C overnight. The cells were treated with 500 U human IFN α

(Calbiochem) for three hours. RNAs were purified from the treated lymphocytes. The Mx1 gene was amplified by RT-PCR using the RNAs as templates and the primer pair AttB1mxF (5'-GGGGACAAGTTTGTACAAAAAAGCAGGCTGTACAGCGTCAAAGAAAAGGAAG-

20 3'/SEQ ID NO: 7) and attB2mxR (5'-GGGGACCACTTTGTACAAGAAAGCTGGGTCTTCTATGATGCTATGCGG-3'/SEQ ID NO: 8). The PCR products were combined with the pDONR201 vector from Gateway Cloning Technology (GIBCO). The mixtures were treated with BP Clonase (GIBCO) at 25°C for one hour. The BP reaction solution was combined with DH5 α competent cells for transformation in a routine fashion, and selection of transformants was performed using LB plates containing 50 μ g/ml kanamycin. The nucleotide sequence of the insert in each clone was determined by direct sequencing of PCR products obtained by colony PCR using attB1mx-F and attB2mx-R (2 minutes at 94°C; 30 cycles of 30 seconds at 94°C, 30 seconds at 64°C, and 3 minutes at 72°C; 5 minutes at 72°C). Normal Mx1 cDNA clone is referred to as pEntmx1, Meishan cDNA clone is referred to as pEntmx1-3, and Landrace cDNA clone is referred to as pEntmx1-11.

[0067]

[Example 2] Construction of Mx1 gene expression vectors

Each of the recombinant clones, pEntmx1, pEntmx1-3, and pEntmx1-11, was inoculated into 15 ml of LB, cultured while shaking at 37°C overnight, and the plasmid DNAs were purified by the alkaline isolation method. The plasmid DNAs were combined with pDEST12.2 (GIBCO). After treatment with LR Clonase (GIBCO) at 25°C for 60 minutes, the DNAs were

transformed into DH5 α competent cells. The selection of transformants was performed using LB containing 100 μ g/m ampicillin. Plasmid DNAs were purified from the respective recombinant clones, which are designated as pExmx1, pExmx1-3, and pExmx1-11. The nucleotide sequence of the insert in each clone was determined by direct sequencing of PCR products obtained by colony PCR using attB1mx-F and attBmx-R.

[0068]

[Example 3] Preparation of Mx1 gene transformants

NIH3T3 cells were passaged and maintained using Dulbecco's modified MEM (DMEM, GIBCO) containing 7% FBS. 5 μ g of pExmx1, pExmx1-3, and pExmx1-11 was each combined with 0.4 ml of 3T3 cell suspension (4×10^6 cells), placed in 2-mm-thick cuvettes, and electroporated at 100 μ F, 13 ohm, 200 v, and 1 kv/cm. After incubation for four days, the cells were cultured in DMEM containing 500 μ g/ml G418 (Geneticin; GIBCO) for the selection of transformants. The pExmx1 transformant is designated as 3T3-0, pExmx1-3 transformant as 3T3-3, and pExmx1-11 transformant as 3T3-11.

[0069]

[Example 4] Preparation of influenza virus particles

An influenza virus (Aichi; H2N2) was inoculated into the allantoic cavities of 11-day-old SPF eggs. The eggs were incubated at 37°C for two days, and then allowed to stand overnight at 4°C. Allantoic fluids were collected from the eggs, and then centrifuged at 3500 rpm for 20 minutes. Aliquots of the supernatant (seed virus solution) were stored at -80°C. The titer of seed virus was estimated to be 107.5 EID₅₀ (50% Egg-Infective Dose)/0.2 ml.

[0070]

[Example 5] Influenza virus infection experiments

Whether forced expression of each Mx1 gene type has an impact on the viral propagation was examined by varying the virus dose used for infection.

[0071]

3T3, 3T3-0, 3T3-3, and 3T3-11 cells were prepared at a cell density of 5×10^4 cells/ml. 1 ml of each cell suspension was added to the wells of a 24-well plate. The cells were incubated in a CO₂ incubator at 37°C for two days, and washed once with phosphate buffered saline (PBS). The seed virus solution was diluted 1:10 (corresponding to MOI (multiplicity of infection)=10) and 1:100 (corresponding to MOI=1) in PBS. The cells were inoculated with the diluted viral solutions, and incubated for adsorption at 37°C for one hour. Unadsorbed virus was removed by washing the cells twice with PBS. 2 ml of DMEM was added to the cells, and the cells were then incubated. Sampling was carried out at 0, 6, 12, 18, 24, 30, 36, 48, and 54 hours. The collected culture supernatants were centrifuged at 2500 rpm for 5 minutes. The supernatants were stored at -80°C until they were assayed for viral titer.

[0072]

Viral titers in the supernatants collected over time were determined as described below. Serial 1:10 dilutions of each viral solution were prepared in PBS. 0.2 ml of each diluted viral solution was inoculated into the allantoic cavities of two 11-day-old eggs. After incubation for two days, the eggs were allowed to stand at 4°C overnight. The allantoic fluids were analyzed for hemagglutinating activity.

[0073]

At MOI=1, the release of propagating virus into culture supernatants was recorded 18 hours after virus infection, and reached its peak of 2.8-3.0 EID₅₀/0.2 ml in 36 hours in the case of 3T3 and 3T3-11 cells. On the other hand, 3T3-0 and 3T3-3 cells have 1.0 EID₅₀/0.2 ml of propagating virus after 36-48 hours. However, no further increase in the viral propagation was recorded. Thus, the virus propagation was clearly different from that observed with the 3T3 and 3T3-11 cells (Fig. 2).

[0074]

In the case of the 3T3 and 3T3-11 cells, at MOI=10, 0.3-0.5 EID₅₀/0.2 ml of infectious virus was detected in culture supernatants 12 hours after virus infection, and the viral titer reached its peak of 4.5-4.7 EID₅₀/0.2 ml in 36 hours. On the other hand, when 3T3-0 and 3T3-3 cells were used, the titer of infectious virus was 0.5 EID₅₀/0.2 ml after 24 hours, 2.7-2.8 EID₅₀/0.2 ml after 48 hours, and no further increase in the viral propagation was recorded after that. Thus, the viral propagation was evidently suppressed (Fig. 3).

[0075]

Influenza virus is an RNA virus. Therefore, there is a possibility that the infection of 3T3 cells may result in the production of mouse IFN α , which affects viral propagation. Accordingly, another infection experiment was carried out, in which cells of each 3T3 cell type were infected with the virus at MOI=10, passaged and maintained in culture media containing anti-mouse IFN α for 12 days. Sampling was carried out after 36 hours. The viral titers in the culture supernatants were determined, and the results showed that the degree of viral propagation was comparable to that with cell groups that had not been treated with anti-mouse IFN α . This suggests that the factor affecting viral propagation is the introduced swine Mx1 gene.

[0076]

[Example 6] Comparison of Mx1 expression levels in transformed cells

To examine whether the difference in viral propagation reflects the level of Mx1 expression in the respective types of cells, RT-PCR was performed using RNAs extracted from the respective types of cells used in the infection experiments.

[0077]

3T3, 3T3-3, and 3T3-11 cells were prepared at a cell density of 2×10^6 cells/ml. RNAs

were extracted and purified from the cells, and the Mx1 expression levels were determined by RT-PCR. The primer pair AttB1mxF and AttB2mxR was used. The PCR conditions were: 30 minutes at 55°C; 2 minutes at 94°C; 40 cycles of 15 seconds at 94°C, 30 seconds at 58°C, and 3 minutes at 68°C; 5 minutes at 72°C for extension. For external standard quantification, mouse G3PDH (TOYOBO) was amplified and used as a positive control for the expression level. A reaction solution to which distilled water was added instead of the template DNA was used as a negative control.

[0078]

According to the results obtained, while the swine Mx1 gene was not expressed in 3T3 cells (lane 1 in Fig. 4), the expression of the swine Mx1 gene was detectable in 3T3-0, 3T3-11, and 3T3-3 cells (lanes 3, 4 and 5 in Fig. 4), and slightly higher in 3T3-11 cells. The expression level of the internal standard mouse G3PDH suggested that the numbers of collected cells did not differ substantially among the respective cell types, and thus the levels of the Mx1 protein produced in 3T3-0, 3T3-11, and 3T3-3 cells were comparable to each other.

[0079]

[Example 7] Detection of an 11-base deletion in the last exon of swine Mx1 gene by PCR amplification

The following primer set was used in PCR for the subject pigs. The portion to be amplified was at positions 1981 to 2160 in the sequence of PubMed accession No. M65087 (Fig. 5).

Primer F: 5'- AGT GAC AGG AGC GAC AAG AG -3' (SEQ ID NO: 9)

Primer R: 5'- CCT GGA GAG TCC GGT TCA -3' (SEQ ID NO: 10)

[0080]

PCR conditions were: 1) 10 minutes at 94°C; 2) 30 seconds at 94°C, 30 seconds at 60°C, and 1 minute at 72°C; 3) 5 minutes at 72°C. Then, 2 µl of the PCR products was electrophoresed on a 6 % polyacrylamide gel at 200 V for 1 hour. The gel was silver stained (Fig. 6). The wild type PCR product had a size of 105 bp. The 11-bp deletion product has a size of 94 bp. The size marker used was fX174/HaeIII.

[0081]

As seen in Fig. 6, the 11-bp deletion could be detected by PCR using the above-described primer pair.

[Effects of the Invention]

[0082]

Among domestic pigs used worldwide, there are some that carry an Mx1 gene containing a 11-bp deletion which results in the loss of the ability to suppress viral propagation.

Such animals are expected to have a serious defect in their defense ability during the early phase of an RNA virus invasion.

[0083]

[Simplified procedures for determining genetic disease resistance]

5 Animals carrying the wild-type gene which confers virus-suppressing ability can be selected from such pig populations, by collecting a small amount of DNA-containing tissues such as flesh and hair root, or blood, and using the extracted DNA to determine genotypes of the Mx1 gene, which is the subject of interest in the present invention. Alternatively, animals comprising undesired genotypes can be eliminated.

10 [0084]

[Improved pig production]

 The level of genetic resistance to diseases caused by RNA viruses, including influenza viruses, can be studied according to the present invention. Healthy animals which are more favorable for pig production can be selected based on the information. Furthermore, the
15 incidence of respiratory diseases in piglets can be reduced, leading to increases in the survival and growth rates.

[0085]

[For human health]

20 Pig populations with impaired ability to suppress influenza virus propagation are susceptible to influenza infection. Unless all of the pigs die, the influenza virus will propagate and undergo genetic mutations, and in some cases evolve into new strains of influenza virus. Like the influenza virus that caused the Spanish influenza epidemic between 1917-1918, resulting in more than 40 million deaths around the world and more than 300,000 deaths in Japan alone, a new strain of influenza virus can cause enormous damage if it infects humans. By
25 selecting pigs with a high ability to suppress influenza virus propagation according to the present invention, the propagation of an influenza virus can be suppressed in pig populations, thereby reducing the chance of the emergence of new influenza virus strains, and raising the possibility of eliminating one threat against humans.

[0086]

[Sequence Listing]

<110> National Institute of Agrobiological Sciences

5 <120> Method to estimate the inherited resistance and susceptibility to RNA virus disease in pigs

<130> MOA-A0214

<140>

10 <141>

<160> 10

<170> PatentIn Ver. 2.1

15

<210> 1

<211> 2545

<212> DNA

<213> Sus scrofa

20

<220>

<221> CDS

<222> (101)..(2092)

25

<220>

<221> polyA_signal

<222> (2517)..(2522)

<400> 1

30 gtaagtgtgg gagaacagcc ctgcatttct gctgacgggt caacgtcaca gcgtcaaaga 60

aaaggaaggt acatttcagc tgaactgac aaggaggaag atg gtt tat tcc agc 115

Met Val Tyr Ser Ser

1

5

35

tgt gaa agt aaa gaa cct gat tca gtt tct gca tcc aat cac ctg tta 163

Cys Glu Ser Lys Glu Pro Asp Ser Val Ser Ala Ser Asn His Leu Leu
 10 15 20
 cta aat ggg aat gat gaa ttg gtg gag aaa agt cac aaa aca ggg cct 211
 5 Leu Asn Gly Asn Asp Glu Leu Val Glu Lys Ser His Lys Thr Gly Pro
 25 30 35
 gag aac aac ctg tac agc cag tac gag gag aaa gtg cgg ccc tgc atc 259
 Glu Asn Asn Leu Tyr Ser Gln Tyr Glu Glu Lys Val Arg Pro Cys Ile
 10 40 45 50
 gac ctg atc gac tca ctg cgg gcc ctg ggc gtg gag cag gac ctg gcc 307
 Asp Leu Ile Asp Ser Leu Arg Ala Leu Gly Val Glu Gln Asp Leu Ala
 55 60 65
 15 ctg ccc gcc atc gcc gtc atc ggg gac cag agt tgc ggc aag agc tcc 355
 Leu Pro Ala Ile Ala Val Ile Gly Asp Gln Ser Ser Gly Lys Ser Ser
 70 75 80 85
 20 gtg ctg gag gcc ctg tgc ggg gtc gct ctg ccc aga ggc agc gga att 403
 Val Leu Glu Ala Leu Ser Gly Val Ala Leu Pro Arg Gly Ser Gly Ile
 90 95 100
 gtg aca aga tgc cct ctt gtg ctg aaa ttg aaa aaa ctg gtg aac gaa 451
 25 Val Thr Arg Cys Pro Leu Val Leu Lys Leu Lys Lys Leu Val Asn Glu
 105 110 115
 gaa gac gaa tgg aag ggc aaa gtc agt tac cgg gac agc gag att gag 499
 Glu Asp Glu Trp Lys Gly Lys Val Ser Tyr Arg Asp Ser Glu Ile Glu
 30 120 125 130
 ctt tca gat gct tgc cag gtg gaa aag gaa gtc agc gca gcc cag att 547
 Leu Ser Asp Ala Ser Gln Val Glu Lys Glu Val Ser Ala Ala Gln Ile
 135 140 145
 35 gcc atc gct ggg gaa ggc gtg gga atc agt cat gag cta atc agt ctg 595

Ala Ile Ala Gly Glu Gly Val Gly Ile Ser His Glu Leu Ile Ser Leu
 150 155 160 165

gag gtc agc tcc cct cat gtc cca gat ctg acc ctc ata gac ctt cct 643
 5 Glu Val Ser Ser Pro His Val Pro Asp Leu Thr Leu Ile Asp Leu Pro
 170 175 180

ggc atc acc agg gta gct gta ggc aat cag cca tac gac atc gaa tac 691
 10 Gly Ile Thr Arg Val Ala Val Gly Asn Gln Pro Tyr Asp Ile Glu Tyr
 185 190 195

cag atc aag tct ctg atc aag aag tac atc tgt aag cag gag acc atc 739
 Gln Ile Lys Ser Leu Ile Lys Lys Tyr Ile Cys Lys Gln Glu Thr Ile
 200 205 210

15 aac ttg gtg gtg gtc ccc tgt aac gtg gac att gcc acc acg gag gcg 787
 Asn Leu Val Val Val Pro Cys Asn Val Asp Ile Ala Thr Thr Glu Ala
 215 220 225

20 ctg cgc atg gcc cag gag gtg gac ccc gaa gga gac agg acc atc ggg 835
 Leu Arg Met Ala Gln Glu Val Asp Pro Glu Gly Asp Arg Thr Ile Gly
 230 235 240 245

atc ttg acg aag ecg gat ctg gtg gac aaa ggc act gag gac aag ata 883
 25 Ile Leu Thr Lys Pro Asp Leu Val Asp Lys Gly Thr Glu Asp Lys Ile
 250 255 260

gtg gac gtg gcg aga aac ctg gtc ttc cac ctg aag aag ggc tac atg 931
 Val Asp Val Ala Arg Asn Leu Val Phe His Leu Lys Lys Gly Tyr Met
 30 265 270 275

att gtc aag tgc agg ggc cag cag gac atc cag gag cag ctg agc ctg 979
 Ile Val Lys Cys Arg Gly Gln Gln Asp Ile Gln Glu Gln Leu Ser Leu
 280 285 290

35 gcc aag gcc ctg cag aag gag cag gcc ttc ttt gaa aac cac gca cat 1027

	Ala Lys Ala Leu Gln Lys Glu Gln Ala Phe Phe Glu Asn His Ala His	
	295	300 305
	ttc agg gat ctt ctg gag gaa ggg cgg gcc acg atc ccc tgc ctg gca	1075
5	Phe Arg Asp Leu Leu Glu Glu Gly Arg Ala Thr Ile Pro Cys Leu Ala	
	310	315 320 325
	gaa aga ctg acc tct gaa ctc atc atg cac atc tgt aaa act ctg ccc	1123
10	Glu Arg Leu Thr Ser Glu Leu Ile Met His Ile Cys Lys Thr Leu Pro	
	330	335 340
	ctg tta gaa aac caa ata aaa gag agt cac cag aaa ata aca gag gag	1171
	Leu Leu Glu Asn Gln Ile Lys Glu Ser His Gln Lys Ile Thr Glu Glu	
	345	350 355
15	ttc cag aag tat ggc tcc gat att cca gag gat gaa agc ggg aag atg	1219
	Leu Gln Lys Tyr Gly Ser Asp Ile Pro Glu Asp Glu Ser Gly Lys Met	
	360	365 370
20	ttt ttt ctg ata gat aaa atc gat gca ttt aat agt gat atc act gct	1267
	Phe Phe Leu Ile Asp Lys Ile Asp Ala Phe Asn Ser Asp Ile Thr Ala	
	375	380 385
	ttg ata caa gga gag gaa ctg gtg gtg gag tac gag tgt cgg ctg ttt	1315
25	Leu Ile Gln Gly Glu Glu Leu Val Val Glu Tyr Glu Cys Arg Leu Phe	
	390	395 400 405
	acc aag atg cga aat gag ttc tgc aga tgg agt gct gtg gtt gaa aag	1363
	Thr Lys Met Arg Asn Glu Phe Cys Arg Trp Ser Ala Val Val Glu Lys	
30	410	415 420
	aat ttc aaa aat ggt tat gac gcc ata tgt aaa caa atc cag ctc ttc	1411
	Asn Phe Lys Asn Gly Tyr Asp Ala Ile Cys Lys Gln Ile Gln Leu Phe	
	425	430 435
35	gaa aat cag tac agg ggg aga gag tgg cca ggg ttt gtg aat tat aag	1459

	Glu Asn Gln Tyr Arg Gly Arg Glu Leu Pro Gly Phe Val Asn Tyr Lys	
	440	445 450
	aca ttt gaa acc atc att aag aag cag gtc agt gtc ctg gaa gag cca	1507
5	Thr Phe Glu Thr Ile Ile Lys Lys Gln Val Ser Val Leu Glu Glu Pro	
	455	460 465
	gcc gtg gac atg ctg cac aca gtg act gat tta gtc cgg ctc gcc ttc	1555
	Ala Val Asp Met Leu His Thr Val Thr Asp Leu Val Arg Leu Ala Phe	
10	470	475 480 485
	aca gat gtt tca gaa aca aat ttt aat gaa ttt ttc aac ctc cac aga	1603
	Thr Asp Val Ser Glu Thr Asn Phe Asn Glu Phe Phe Asn Leu His Arg	
	490	495 500
15	act gcc aag tcc aaa att gaa gac att aaa tta gaa caa gaa aaa gaa	1651
	Thr Ala Lys Ser Lys Ile Glu Asp Ile Lys Leu Glu Gln Glu Lys Glu	
	505	510 515
	gct gag acg tcg atc cgg ctc cac ttc caa atg gag cag atc gtg tac	1699
20	Ala Glu Thr Ser Ile Arg Leu His Phe Gln Met Glu Gln Ile Val Tyr	
	520	525 530
	tgc cag gac cag gtc tat cgg gcc gcg ctg cag aag gtc aga gag aag	1747
25	Cys Gln Asp Gln Val Tyr Arg Gly Ala Leu Gln Lys Val Arg Glu Lys	
	535	540 545
	gag gcg gaa gaa gaa aag aac aga aaa tca aac cag tac ttt ctg tcg	1795
	Glu Ala Glu Glu Glu Lys Asn Arg Lys Ser Asn Gln Tyr Phe Leu Ser	
30	550	555 560 565
	tcg ccg gcc ccc tcc tca gac ccc tcc ata gcc gag atc ttt cag cac	1843
	Ser Pro Ala Pro Ser Ser Asp Pro Ser Ile Ala Glu Ile Phe Gln His	
	570	575 580
35	ctg att gcc tac cat cag gag gtc gcc aag cgc atc tcc agc cac atc	1891

Leu Ile Ala Tyr His Gln Glu Val Gly Lys Arg Ile Ser Ser His Ile
 585 590 595

5 cct ctg atc atc cag ttc ttc atc etc cgg acc ttt ggg cag cag ctg 1939
 Pro Leu Ile Ile Gln Phe Phe Ile Leu Arg Thr Phe Gly Gln Gln Leu
 600 605 610

10 cag aag agc atg ctg cag ctg ctg cag aac aag gac caa tac gac tgg 1987
 Gln Lys Ser Met Leu Gln Leu Leu Gln Asn Lys Asp Gln Tyr Asp Trp
 615 620 625

ctc ctg agg gag cgc agt gac acc agc gac aag agg aag ttc ctg aag 2035
 Leu Leu Arg Glu Arg Ser Asp Thr Ser Asp Lys Arg Lys Phe Leu Lys
 630 635 640 645

15 gag cgg ctg atg cgg ctg acc cag gct cgg cgc cgg etc gcc aag ttc 2083
 Glu Arg Leu Met Arg Leu Thr Gln Ala Arg Arg Arg Leu Ala Lys Phe
 650 655 660

20 cca ggc tga accggactct ccaggcgccc cggggctctcc agggcacgtc 2132
 Pro Gly
 tccaggcaac gaggaccaac ctccctccct aacagactag catcatgagc tctgtttcg 2192

25 cacatctctc tgtggttagt agactctaaa gccaccgtcc ctgctgttag ttgctgagga 2252
 cttagcaaga agctgtgata agcacgtcgg ctgcaagcat caggccattt acttgaatga 2312
 gccccgccaa cgcttcgcct cccgcgcctc tctccatccc tctctccatc cttctctcca 2372

30 tcctgtgata ggatactggt cccgcgatag catcatagaa gggtcattct ggtttctgta 2432
 caagcctttc acgcccaatg tcttaggggc attacagcca cctgtgtgga tggatgcaca 2492

35 tagaagccta tttcttttat ttgtaataaa ctgggttcta ccagcaaaaa aaa 2545

<210> 2

<211> 663

<212> PRT

5 <213> Sus scrofa

<400> 2

Met Val Tyr Ser Ser Cys Glu Ser Lys Glu Pro Asp Ser Val Ser Ala

1 5 10 15

10

Ser Asn His Leu Leu Leu Asn Gly Asn Asp Glu Leu Val Glu Lys Ser

20 25 30

His Lys Thr Gly Pro Glu Asn Asn Leu Tyr Ser Gln Tyr Glu Glu Lys

15 35 40 45

Val Arg Pro Cys Ile Asp Leu Ile Asp Ser Leu Arg Ala Leu Gly Val

50 55 60

20 Glu Gln Asp Leu Ala Leu Pro Ala Ile Ala Val Ile Gly Asp Gln Ser

65 70 75 80

Ser Gly Lys Ser Ser Val Leu Glu Ala Leu Ser Gly Val Ala Leu Pro

85 90 95

25

Arg Gly Ser Gly Ile Val Thr Arg Cys Pro Leu Val Leu Lys Leu Lys

100 105 110

Lys Leu Val Asn Glu Glu Asp Glu Trp Lys Gly Lys Val Ser Tyr Arg

30 115 120 125

Asp Ser Glu Ile Glu Leu Ser Asp Ala Ser Gln Val Glu Lys Glu Val

130 135 140

35 Ser Ala Ala Gln Ile Ala Ile Ala Gly Glu Gly Val Gly Ile Ser His

145 150 155 160

Glu Leu Ile Ser Leu Glu Val Ser Ser Pro His Val Pro Asp Leu Thr
 165 170 175

5 Leu Ile Asp Leu Pro Gly Ile Thr Arg Val Ala Val Gly Asn Gln Pro
 180 185 190

Tyr Asp Ile Glu Tyr Gln Ile Lys Ser Leu Ile Lys Lys Tyr Ile Cys
 195 200 205

10 Lys Gln Glu Thr Ile Asn Leu Val Val Val Pro Cys Asn Val Asp Ile
 210 215 220

Ala Thr Thr Glu Ala Leu Arg Met Ala Gln Glu Val Asp Pro Glu Gly
 15 225 230 235 240

Asp Arg Thr Ile Gly Ile Leu Thr Lys Pro Asp Leu Val Asp Lys Gly
 245 250 255

20 Thr Glu Asp Lys Ile Val Asp Val Ala Arg Asn Leu Val Phe His Leu
 260 265 270

Lys Lys Gly Tyr Met Ile Val Lys Cys Arg Gly Gln Gln Asp Ile Gln
 275 280 285

25 Glu Gln Leu Ser Leu Ala Lys Ala Leu Gln Lys Glu Gln Ala Phe Phe
 290 295 300

Glu Asn His Ala His Phe Arg Asp Leu Leu Glu Glu Gly Arg Ala Thr
 30 305 310 315 320

Ile Pro Cys Leu Ala Glu Arg Leu Thr Ser Glu Leu Ile Met His Ile
 325 330 335

35 Cys Lys Thr Leu Pro Leu Leu Glu Asn Gln Ile Lys Glu Ser His Gln
 340 345 350

Lys Ile Thr Glu Glu Leu Gln Lys Tyr Gly Ser Asp Ile Pro Glu Asp
 355 360 365

5 Glu Ser Gly Lys Met Phe Phe Leu Ile Asp Lys Ile Asp Ala Phe Asn
 370 375 380

Ser Asp Ile Thr Ala Leu Ile Gln Gly Glu Glu Leu Val Val Glu Tyr
 385 390 395 400

10 Glu Cys Arg Leu Phe Thr Lys Met Arg Asn Glu Phe Cys Arg Trp Ser
 405 410 415

Ala Val Val Glu Lys Asn Phe Lys Asn Gly Tyr Asp Ala Ile Cys Lys
 15 420 425 430

Gln Ile Gln Leu Phe Glu Asn Gln Tyr Arg Gly Arg Glu Leu Pro Gly
 435 440 445

20 Phe Val Asn Tyr Lys Thr Phe Glu Thr Ile Ile Lys Lys Gln Val Ser
 450 455 460

Val Leu Glu Glu Pro Ala Val Asp Met Leu His Thr Val Thr Asp Leu
 465 470 475 480

25 Val Arg Leu Ala Phe Thr Asp Val Ser Glu Thr Asn Phe Asn Glu Phe
 485 490 495

Phe Asn Leu His Arg Thr Ala Lys Ser Lys Ile Glu Asp Ile Lys Leu
 30 500 505 510

Glu Gln Glu Lys Glu Ala Glu Thr Ser Ile Arg Leu His Phe Gln Met
 515 520 525

35 Glu Gln Ile Val Tyr Cys Gln Asp Gln Val Tyr Arg Gly Ala Leu Gln
 530 535 540

Lys Val Arg Glu Lys Glu Ala Glu Glu Glu Lys Asn Arg Lys Ser Asn
 545 550 555 560

5 Gln Tyr Phe Leu Ser Ser Pro Ala Pro Ser Ser Asp Pro Ser Ile Ala
 565 570 575

Glu Ile Phe Gln His Leu Ile Ala Tyr His Gln Glu Val Gly Lys Arg
 580 585 590

10

Ile Ser Ser His Ile Pro Leu Ile Ile Gln Phe Phe Ile Leu Arg Thr
 595 600 605

15

Phe Gly Gln Gln Leu Gln Lys Ser Met Leu Gln Leu Leu Gln Asn Lys
 610 615 620

Asp Gln Tyr Asp Trp Leu Leu Arg Glu Arg Ser Asp Thr Ser Asp Lys
 625 630 635 640

20

Arg Lys Phe Leu Lys Glu Arg Leu Met Arg Leu Thr Gln Ala Arg Arg
 645 650 655

Arg Leu Ala Lys Phe Pro Gly
 660

25

<210> 3

<211> 23

<212> DNA

30

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:an artificially
 synthesized primer sequence

35

<400> 3

ctgaaagatc tcggctaagg agg

23

<210> 4

5 <211> 25

<212> DNA

<213> Artificial Sequence

<220>

10 <223> Description of Artificial Sequence:an artificially
synthesized primer sequence

<400> 4

aagaagctga gacgtcgatc cggct

25

15

<210> 5

<211> 21

<212> DNA

20 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:an artificially
synthesized primer sequence

25

<400> 5

aagcgcatct ccagccacat c

21

30 <210> 6

<211> 20

<212> DNA

<213> Artificial Sequence

35 <220>

<223> Description of Artificial Sequence:an artificially

synthesized primer sequence

<400> 6

aagacattgg gcgtgaaagg

20

5

<210> 7

<211> 53

<212> DNA

10 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:an artificially
synthesized primer sequence

15

<400> 7

ggggacaagt ttgtacaaa aagcaggctg tcacagcgtc aaagaaaagg aag

53

20 <210> 8

<211> 49

<212> DNA

<213> Artificial Sequence

25 <220>

<223> Description of Artificial Sequence:an artificially
synthesized primer sequence

<400> 8

30 ggggaccact ttgtacaaga aagctgggtc ctctatgat gctatgagg

49

<210> 9

<211> 20

35 <212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:an artificially
synthesized primer sequence

5

<400> 9

agtgacagga gcgacaagag

20

10 <210> 10

<211> 18

<212> DNA

<213> Artificial Sequence

15 <220>

<223> Description of Artificial Sequence:an artificially
synthesized primer sequence

<400> 10

20 cctggagagt ccggttca

18

[Brief Description of the Drawings]

[Fig. 1] A photograph showing a three-way crossbred pig used for fattening/production.

[Fig. 2] A graph showing the ability of each Mx1 genotype to suppress viral propagation at an MOI of 1.

5 [Fig. 3] A graph showing the ability of each Mx1 genotype to suppress viral propagation at an MOI of 10.


[Fig. 4] A photograph that compares the Mx1 expression levels in the respective transformed cells of 3T3, 3T3-0, 3T3-3, and 3T3-11, based on the results obtained using RT-PCR.

10 [Fig. 5] A figure showing an "11-bp deletion" region in the last exon of the swine Mx1 gene, and nucleotide sequences of a primer pair to be used for amplifying the region.


[Fig. 6] A photograph showing the detection of an 11-base deletion in the last exon of the swine Mx1 gene using PCR amplification.

[Document Name] Drawings

[Fig. 1]




[Fig. 2]




5

[Fig. 3]




[Fig. 4]



[Fig. 5]

CGACTGGCTC	CTGAGGGAGC	<u>GCAGTGACAC</u>	<u>CAGCGACAAG</u>	<u>AGGAAGTTC</u>	TGAAGGAGCG
PRIMER F					
GCTGATGCGG	CTGACCCAGG	<u>CTCGGCGCCG</u>	<u>GCTCGCCAAG</u>	<u>TTCCAGGCT</u>	<u>GAACCGGACT</u>
11-bp DELETION					
PRIMER R					
<u>CTCCAGGCGG</u>	<u>CCGGGGGTCT</u>	<u>CCAGGGCAGG</u>	<u>TCTCCAGGCA</u>	<u>ACGAGGACCA</u>	<u>ACCTCCTTCC</u>

5 [Fig. 6]



[Document Name] Abstract

[Abstract]

[Problems to Be Solved]

- 5 An objective of the invention is to provide methods for determining genetic resistance of pigs to diseases caused by RNA viruses.

[Means for Solving the Problems]

- 10 The inventors investigated the impact of an 11-bp deletion in the swine Mx1 gene on the ability to suppress propagation of influenza viruses belonging to the myxovirus family, and revealed that the deletion led to a complete loss of the ability to suppress viral propagation.
- 10 Through the detection of the 11-bp deletion, pigs can be examined for their resistance to influenza viruses.

[Selected Drawings] None